

## Changes in Monocyte Populations Following Acute Aerobic Exercise in Breast Cancer Survivors

**Khosravi N:** Physical Education & Sport Sciences Department, Faculty of Humanities, Tarbiat Modares University, Tehran, Iran & Department of Exercise & Sport Science, Exercise Oncology Research Laboratory, University of North Carolina, Chapel Hill, NC USA

**Hanson ED:** Department of Exercise & Sport Science, Exercise Oncology Research Laboratory, University of North Carolina, Chapel Hill, NC USA

**Farajivafa V:** Physical Education & Sport Sciences Department, Faculty of Humanities, Tarbiat Modares University, Tehran, Iran & Department of Exercise & Sport Science, Exercise Oncology Research Laboratory, University of North Carolina, Chapel Hill, NC USA

**Agha-Alinejad H:** Physical Education & Sport Sciences Department, Faculty of Humanities, Tarbiat Modares University, Tehran, Iran

**Haghighat S:** Quality of Life Department, Breast Cancer Research Center, Motamed Cancer Institute, ACECR, Tehran, Iran

**Molannouri Shamsi M:** Physical Education & Sport Sciences Department, Faculty of Humanities, Tarbiat Modares University, Tehran, Iran

**Evans WS, Lee JT, Danson E, Wagoner CW, Harrell EP:** Department of Exercise & Sport Science, Exercise Oncology Research Laboratory, University of North Carolina, Chapel Hill, NC USA

**Nyrop KA, Muss HB:** Department of Hematology Oncology University of North Carolina, Chapel Hill, NC USA

**Bartlett DB:** Department of Medicine, Duke University, Durham NC, USA

**Battaglini CL:** Department of Exercise & Sport Science, Exercise Oncology Research Laboratory, University of North Carolina, Chapel Hill, NC USA

**Corresponding Author:** Hamid Agha-Alinejad, halinejad@modares.ac.ir

### Abstract

**Introduction:** Exercise is now strongly recommended for breast cancer patients to improve their overall health and quality of life. Monocytes play an important role in the cancer immune system and a better understanding of how acute exercise alters the monocyte subsets would aid in exercise prescription.

**Methods:** Ten breast cancer survivors (age:  $59 \pm 7.1$ ) who completed their primary cancer treatment within the previous year were evaluated in this study. Using flow cytometry, monocyte subset percentages were evaluated before, immediately after, and 1 hour after 45 minutes of acute, intermittent exercise. Exercise intensity was 60% of peak wattage obtained from a cardiopulmonary exercise test.

**Results:** The percentage of CD14<sup>+</sup> monocytes and CD14<sup>+</sup>CD16<sup>-</sup> monocyte subsets changed significantly across the trial ( $p = 0.016$  and  $p = 0.016$ , respectively), with a small, non-significant increase immediately after exercise (CD14<sup>+</sup>: 9%,  $p = 0.314$ ; CD14<sup>+</sup>CD16<sup>-</sup>: 5%,  $p = 0.594$ ) followed by a larger significant decrease 1 hour after exercise relative to baseline (CD14<sup>+</sup>: -26%,  $p = 0.015$ ; CD14<sup>+</sup>CD16<sup>-</sup>: -28%,  $p = 0.021$ ). CD14<sup>+</sup>CD16<sup>+</sup> subpopulation showed a tendency to change across the trial but this did not quite reach significance ( $p = 0.097$ ).

**Conclusion:** These findings suggest that acute intermittent exercise mobilizes CD14<sup>+</sup> monocytes and CD14<sup>+</sup>CD16<sup>-</sup> monocyte subsets in breast cancer survivors in a manner that is comparable to previous reports in healthy individuals. Further studies are warranted to determine the functionality of the mobilized monocytes and the effects of exercise training.

**Keywords:** Exercise, Breast Cancer, Monocyte, Immune System