

Changes in Monocyte Populations Following Acute Aerobic Exercise in Breast Cancer Survivors

Khosravi N: Physical Education & Sport Sciences Department, Faculty of Humanities, Tarbiat Modares University, Tehran, Iran & Department of Exercise & Sport Science, Exercise Oncology Research Laboratory, University of North Carolina, Chapel Hill, NC USA

Hanson ED: Department of Exercise & Sport Science, Exercise Oncology Research Laboratory, University of North Carolina, Chapel Hill, NC USA

Farajivafa V: Physical Education & Sport Sciences Department, Faculty of Humanities, Tarbiat Modares University, Tehran, Iran & Department of Exercise & Sport Science, Exercise Oncology Research Laboratory, University of North Carolina, Chapel Hill, NC USA

Agha-Alinejad H: Physical Education & Sport Sciences Department, Faculty of Humanities, Tarbiat Modares University, Tehran, Iran

Haghighat S: Quality of Life Department, Breast Cancer Research Center, Motamed Cancer Institute, ACECR, Tehran, Iran

Molannouri Shamsi M: Physical Education & Sport Sciences Department, Faculty of Humanities, Tarbiat Modares University, Tehran, Iran

Evans WS, Lee JT, Danson E, Wagoner CW, Harrell EP: Department of Exercise & Sport Science, Exercise Oncology Research Laboratory, University of North Carolina, Chapel Hill, NC USA

Nyrop KA, Muss HB: Department of Hematology Oncology University of North Carolina, Chapel Hill, NC USA

Bartlett DB: Department of Medicine, Duke University, Durham NC, USA

Battaglini CL: Department of Exercise & Sport Science, Exercise Oncology Research Laboratory, University of North Carolina, Chapel Hill, NC USA

Corresponding Author: Hamid Agha-Alinejad, halinejad@modares.ac.ir

Abstract

Introduction: Exercise is now strongly recommended for breast cancer patients to improve their overall health and quality of life. Monocytes play an important role in the cancer immune system and a better understanding of how acute exercise alters the monocyte subsets would aid in exercise prescription.

Methods: Ten breast cancer survivors (age: 59 ± 7.1) who completed their primary cancer treatment within the previous year were evaluated in this study. Using flow cytometry, monocyte subset percentages were evaluated before, immediately after, and 1 hour after 45 minutes of acute, intermittent exercise. Exercise intensity was 60% of peak wattage obtained from a cardiopulmonary exercise test.

Results: The percentage of $CD14^+$ monocytes and $CD14^+CD16^-$ monocyte subsets changed significantly across the trial ($p = 0.016$ and $p = 0.016$, respectively), with a small, non-significant increase immediately after exercise ($CD14^+$: 9%, $p = 0.314$; $CD14^+CD16^-$: 5%, $p = 0.594$) followed by a larger significant decrease 1 hour after exercise relative to baseline ($CD14^+$: -26%, $p = 0.015$; $CD14^+CD16^-$: -28%, $p = 0.021$). $CD14^+CD16^+$ subpopulation showed a tendency to change across the trial but this did not quite reach significance ($p = 0.097$).

Conclusion: These findings suggest that acute intermittent exercise mobilizes $CD14^+$ monocytes and $CD14^+CD16^-$ monocyte subsets in breast cancer survivors in a manner that is comparable to previous reports in healthy individuals. Further studies are warranted to determine the functionality of the mobilized monocytes and the effects of exercise training.

Keywords: Exercise, Breast Cancer, Monocyte, Immune System